



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

302. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Prove that the system of equations

$$xu - 5yv = 2,$$

$$xv + yu = 1,$$

has no integral solution in x, y, u, v except those for which one of the unknowns is zero.

GEOMETRY.

335. Proposed by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Determine analytically, the point where three lines in a plane appear of equal length.

CALCULUS.

260. Proposed by V. M. SPUNAR, East Pittsburg, Pa.

A natural equation of a surface may be defined as an equation in which the differential quotients of the principal radius, ρ , of curvature to the element of arc in the direction of the principal curvature are shown as a function of ρ , $\frac{d^n \rho}{ds} = F(\rho)$. Required the natural equation of the whole surface of second power.

261. Proposed by S. A. COREY, Hiteman, Iowa.

Prove that $\sum_{x=1}^{x=\infty} \frac{1}{a + 2bx^2 + cx^4} = \frac{\pi}{\sqrt{[8ac(\sqrt{ac+b})]}} - \frac{1}{2a}$, where $ac > b^2$.

262. Proposed by H. SCHAFFER, Fayetteville, Ark.

Prove that the circle is the only plane curve of constant curvature.

MECHANICS.

217. Proposed by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Given, the mean distance from earth to sun, 1.49×10^{15} centimeters; radius of the earth, 6.37×10^8 centimeters; velocity of the earth in its orbit, 2.96×10^6 centimeters per second; velocity of rotation of a point on the equator, 4.63×10^4 centimeters per second; mass of the earth, 6.14×10^{27} grams; find (1) the total energy of the earth in ergs; (2) the angular velocity of the earth on its axis; and (3) the angular velocity of the earth around the sun.

218. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Cut a uniform, circular cylinder by two planes whose line of intersection is without the cylinder. The centroid G of the surface of the portion of the cylinder thus cut off lies in a plane elliptic section, in which plane also lies the line of intersection aforesaid. C is the center of the ellipse, and the pole of the intersection line with reference to this ellipse is X . Show (1) that C, X , and G are collinear, and (2) that $XC = 2CG$.